

New Meteorological and Geological Study of Acquapendente (VT)

Tasselli, D.

TS Corporation Srl - Astronomy and Astrophysics Department
Regione Salaria, 10010 Andrate TO - Italy
E-mail:diego.tasselli@tscorporation.org

Ricci, S.

TS Corporation Srl - Meteorological and Climatic Change Department
Regione Salaria, 10010 Andrate TO - Italy
E-mail:stefano.ricci@tscorporation.org

Bianchi, P.

TS Corporation Srl - Geology and Geophysics Department
Regione Salaria, 10010 Andrate TO - Italy
E-mail:pamela.bianchi@tscorporation.org

Abstract

In this paper we present an analysis of the geological, meteorological and climatic data recorded in Acquapendente (VT) over 24 years. These data are compared to check local variations, long term trends, and correlation with mean annual temperature. The ultimate goal of this work is to understand long term climatic changes in this geographic area. The analysis is performed using a statistical approach. From each long series of data calculate the hourly averages and that the monthly averages in order to reduce the fluctuations in the short time due to the day / night cycle. A particular care is used to minimize any effect due to prejudices in case of lack of data. Finally, we calculate the annual average from the monthly ones.

Keyword: atmospheric effects - site testing - earthquake data - geological model - methods: statistical.

This paper was prepared with the $\text{\LaTeX}$

1 Introduction

In this paper we present an analysis of geological, meteorological and climatic data released in Acquapendente (VT) by “SMCS - Stazione Meteo-Climatica e Sismologica” a project by Meteorological and Climatic Change and Geological Depart-

ment of TS Corporation Srl.[1].

We present for the first time an analysis of measurements obtained from local geological [11] and meteorological data [4][5] and compared in order to check local variations or meteorological conditions. We discuss the geological data, annual temperature means and differences between day

time and night time mean values and their comparison with the down time.

1.1 Location

Acquapendente is a little town in the center of Italy. Here are identifiable altimeter data, the geographic coordinates and seismic data.

Latitudine	Longitude
42°44'41,28" N	11°51'54,36" E

Geographic Data of Acquapendente

Acquapendente is located in the extreme north of Latium region near the border with Tuscany region, whose boundaries are marked by a sudden from San Casciano lake, and Umbria region, about six miles north of Bolsena lake at the Natural Reserve of Rufeno's Mount that located on the opposite side of the river Paglia; the town is crossed by the Cassia's street. Acquapendente owes its name to the fact that it is located near several small waterfalls that flow into the river Paglia. The geological map of the town is shown in Geological and Seismic data section. [3]

Seismic Zone	Description
2B	Area with average seismic hazard may occur where earthquakes strong enough. The sub area 2B indicates a value of $ag < 0,20g$.

Seismic Data of Acquapendente

2 Annual data analysis

Summers are generally warm and winters are fairly rigid and often winds blow from the north that may last a long time. In winter mornings it is easy to find fog also due to the proximity of the Paglia river. The snow appears about two or three times a year, sometimes in large quantities. The rains dominate the autumn and spring seasons. The following table identifies climate data assigned by Decree of the President of the Republic n. 412 of 26 August 1993.[2]. Characterized by a fairly humid climate,

the summer can often record the maximum temperature higher than the district, while in winter with thermal inversion and the particular conditions in which it occurs, it manages to be the coldest city of area, and among the coldest in central Italy (of course compared to the cities in which it is carried out monitoring data). It has a climate for "extreme" in both summer and winter conditions like temperature range and inversion are the order of the day.

2.1 The temperature inversion

The temperature inversion, as the name implies, is when the vertical thermal gradient is reversed, ie, when the temperature increases with altitude (positive gradient) instead of decreasing (negative gradient). Basic conditions so that we can experience nocturnal inversion are essentially three:

- Preparation of morphological country to this phenomenon;
- Terms of persistent high pressure (skies);
- Lack of wind.

The summation of these factors causes a marked loss of heat due to radiation night which leads to the creation of a bearing of cold air that station in the lower layers.

This means that the bearing in the early morning minimum Acquapendente registers much lower than the neighboring countries with higher altitude.

Climatic Zone	Day Degrees
E	2,299

Climatic Parameter of Acquapendente

3 Meteo-Climatic Parameter

In this section we describe air temperatures (T), Dew Point, Humidity, Pressure, Day Time and Night Time Variation, Rain's Days and Fog's Day, obtained by an accurate analysis of the meteorological data from local data by archive [4]. We analyzed the parameters given in Table 1. Should be noted that the values considered

are related to the last twenty-four-year average and made available for the period 1990-2014.

Average Annual Temperature	14,33°C
T average warmest (ago-03)	28,27°C
T average coldest (feb-12)	1,42°C
Annual temperature range	9,97°C
Months with average T > 20°C	83
Total rainfall 1990-2014 [mm]	21401,93
Rain Days	1198
Fog Days	912
Storm Days	181
Rain/Storm Days	506
Rain/Snow Days	21
Rain/Fog Days	159
Rain/Thunder/Fog Days	68
Snow Days	22
Wind Speed max Km/h (Apr-2008)	36,57
Wind Speed min Km/h (Dec-2011)	9,00
Rain max mm (Nov-2008)	285,90
Rain min mm (Jun-2004)	1,50
Earthquake Min (2011/12/20)	0,2 Mw
Earthquake Max (2006/07/06)	3,6 Mw
Earthquake Deep Min (2012/08/21)	0,7 Km
Earthquake Deep Max (2010/04/28)	42,5 Km

Parameter of this Study

3.1 Solar Radiation Territory

The data irradiation of territory taken from the parameters and the data prepared by the European Union, demonstrate the trend of irradiation for the municipality, visible in next table:

Month	DNI	Month	DNI
Jan	2410	Feb	3500
Mar	4030	Apr	4800
May	5570	Jun	6840
Jul	7990	Aug	7080
Sep	5100	Oct	3710
Nov	2620	Dec	2290
Year	4670		


Direct Normal Irradiance (Wh/m²/day)

The weather data and the graphs show extrapolated for the territory covered by the study, including a radiation in the range

between 1200 and 1300 kWh /1kWp as map prepared by the European Union [13] and visible in figure 18, characterized in Over the months, irradiation presented in the graph in Figure 19, which shows the data of the table above, which shows the territory of Acquapendente, a total irradiance Annual of 4670 Wh/m²/day.

3.2 Temperature


In this section we describe air temperatures (T) obtained by an accurate analysis of the meteorological data. The year average temperature has a tendency to remain stable throughout the study period, except: decrease in the period between 1990 and 1996; back and stabilize during 1997-2014. Next graphics show this evidence:



Relationship of Year Temperature and Average 1990-2014.

The values are calculated considering the entire measurement period (1990-2014), drawing on data from the Annuals published by "Il Meteo.it" [4] for the period 1990-2014.

The study period showed an increase in "heat waves" such as that of 2003, whose negative impact is amplified by the increase in temperature in the urban areas, creating the effect of "heat island" mainly due to soil sealing.



Variation of Month Temperature about year 1990-2002 and 2003-2014

The graphs shown in Figures 4 and 5

show the performance of the maximum and minimum temperatures proportional to the average of the period of study (1990-2014). The study shows the following trends:

- Minimum temperatures: there was an increase of the values recorded with a higher peak in February 2012, with 1,42°C;
- Maximum temperatures: evidence for the period 2007-2009 is an increasing trend compared to the 1990-2014 average, while evidence a decrease in the period 1990-1997. The maximum temperature was 28.27°C in August 2003;

Table 5 shows the trend of the points of maximum temperature expected in the summer quarter (June-July-August) and are shown in Figure 15. Study shows that the month of August 2003 recorded the highest average maximum value in a context in which the entire month recorded in the period 1990-2014 values always above average.

Always Table 5 shows the trend of the points of minimum temperature expected in the winter quarter (January-February-March) and are shown in Figure 16. Study shows that the month of February recorded the highest average minimum value, in a context in which the entire month recorded in the period 1990-2014 values always above average. Studying in particular the months with the values of minimum temperature and maximum minors, the following is noted:

- the month of August (characterized by the presence of the highest value observed in maximum temperatures), shows that trends in temperature has been getting consistently below average for the period 1990 to 2014, while there were two exceedances of this value over the years 2003 and 2012, in agreement with what evident from the graph in Figure 13;

- the month of February (characterization from the value of the lowest minimum temperature for the period of study), shows that the trend of temperatures has always been, in agreement with what reported in the graph in Figure 14.

3.3 Dew point

In this section we describe Dew point obtained by an accurate analysis of the meteorological data. This study showed:

- an increase in the year: 1994, 2001, 2002 (the highest Dew Point period for this study), 2004, 2005, 2006, 2007, 2009, 2010, 2011, 2012 and 2014;
- a decrease in rest of study period, with the lowest Dew Point in 1991 year.

Graphics in Figure 11 show this trend.

3.4 Humidity

In this section we describe humidity obtained by an accurate analysis of the meteorological data. The study highlights a gap in the annual humidity values equal to 67,56%, calculated according to this formula:

$$\frac{av}{am} \quad (1)$$

av = annual value, *am* = average moisture 1990-2014

The graph in Figure 9 shows a trend tends to be stable in the values obtained with threshold in growth over the period 1990,1992,1994,1995,1996,2002,2004 and 2010, and a decrease in the period 1991, 1993, 1998,1999, 2000, 2003, 2006, 2007, 2009, 2011 and 2012.

3.5 Pressure

In this section we describe pressure obtained by an accurate analysis of the Pressure data.

The analysis of the data showed:

- an increase in atmospheric pressure over the years: 1991, 1998, 2000 (year with the highest level of pressure throughout the study period), 2003, 2004, 2007, 2008, 2011, 2012 and 2014;
- a decrease in atmospheric pressure over the years: 1993, 1994, 1995, 1996, 1997, 2001, 2002, 2005 (the year with the lowest level of pressure throughout the study period), 2006, 2010 and 2013.

In Figure 10 we can see the Pressure Graphics for this study.

3.6 Day time and night time variation


In this section we describe number of Day time and night time variations obtained by an accurate analysis of the meteorological data. The annual averages of the differences between day time and night time temperatures ΔT have been computed and the results are reported in Table 2. Also in Figure 3, we can see the plot of oscillations of the ΔT seem to reduce the amplitude during the years.

The difference of temperatures, with an average difference of $9,97^{\circ}\text{C}$ (see table 2) does confirm the trend indicate Acquapendente as one of Italian municipalities with higher ratio of the temperature difference between day and night. Table 2 and Figure 3 show these effects.

3.7 Rain's Days

In this section we describe number of Rain's days obtained by an accurate analysis of the meteorological data.

An increase in extreme weather events and abnormal, leading to a potential increase in precipitation intensity for each event, especially in areas where there is an increase of average precipitation.




Days Rain for Month from 1990 to 2014.

The increase in intense rainfall of short duration boosts the risk of flash-food, fast flooding of an area circumscribed morphologically, due to the fast “saturation” of the surface soil that is no longer able to absorb the rain. The episodes of intense precipitation can also determine the phenomena of surface runoff of rainwater with a possible increase in flooding, but also the risk of water pollution (pollutants from agricultural and road runoff).

The study period showed an abnormal increase in rainfall. The pie chart shows the ratio between the total mm of rain measured in a year and the total rainfall measured in the period 1990-2014, according to this formula:


$$TotalRain = \frac{a}{b} \times 100 \quad (2)$$

a = annual rainfall value, **b** = total value of the rain period



% of Rain for this study.

In Figure 6 and 7 we can see the total of rain/year in this study. The values are visible in the table 3. Next pie chart evidence type of rain for this study.



Type of Rain for this study.


Figure 17 evidence the total rain in mm, for only month.

The study shows an abnormal increase in rains, shown in Figure 7, in the years: 2014, 2013, 2012, 2009 (the highest year period mm of rain fell), 2008, 2006, 2005, 2004, 1996, 1995, 1992, with an average increase of 126.73% of the total annual rainfall mm which is equal to 856.08 mm on average for year.

3.8 Fog's days

In this section we describe number of Fog's days obtained by an accurate analysis of the meteorological data.


The study showed a trend increase in the presence of days with fog and evidence that October is the first month for number of Fog's day and July are the laster. Figure 8 we can see the number of Fog days by this study.



Days Fog for Month from 1990 to 2014

4 Geological and Earthquake data

In this section we describe the Geological and Earthquake data.



Geological Map of Acquapendente - VT [3]

The geology in Acquapendente is characterized by high heat flow, locally more than 200 mWm^{-2} [8] positive magnetic anomalies [9], shallow earthquakes, positive gravity anomalies and crustal thinning (20-25 km). The area lies within the catchment area of the river Paglia, and the percentage composition of the various lithologies in the basin is so divided: 37.5% volcanics, flooding 20%, 17.5% flysch, anthropogenic debris cones and 5%, clay 15%, sands 2%.

Acquapendente falls in complex lavas and ignimbrites lithoid, and consists of lava flows and ignimbrites lithoid (Pliocene-Pleistocene) interspersed at various levels in the complex pyroclastic. The thickness of this complex is extremely variable and ranges between a few dozen to over a hundred meters. The permeability of the complex is variable and is related to the different degree of cracking and the presence of tectonic reasons. In relative terms the degree of permeability and to be considered medium to high and, together with the complex pyroclastic constitutes the main aquifer area under examination. The field observations combined with the information of the surveys have allowed to estimate more than 2500 m in thickness of the Pliocene deposits [11].

The predominant lithology consists of bluish clay, interspersed with different stratigraphic levels from debris flow conglomerate and sandstone turbidites. These sediments have been described in detail by [12] belong to the first cycle of sedimentation [11].

Sedimentation during this time period occurred throughout the basin and is also demonstrated by the polls and deep wells for geothermal exploration that have met these sediments beneath the volcanics..

4.1 Description of the main outcrop formations.

The magmatic Pliocene-Pleistocene (rocks apparatus Vulsino) outcropping in the territory of Acquapendente are:

- **Basanites and tephrites leucititiche:** The basanites emerge around Mt. Rosso with a dark gray casting very compact. The tephrites leucititiche are present with extensive lava outcropping around Acquapendente. Those visible along the right side of Straw differ from others because of the abundance of phenocrysts of leucite and alteration that turns into a reddish sandy soil.
- **Leucitites passers tephrites leucititici and basalts:** The leucitites and leucitites passers tephrites rocks are dark gray compact very fine. They are found in lava flows at Acquapendente. The leucitites passers in basalts leucititici alternate, near Mt. Landro, with banks of lapilli and scoria.
- **Tuffs yellow pumice gray:** Rest directly on the flysch and the Pliocene as the lava flows and other types of pyroclastic and reach considerable power that exceeds the 50 m. They are characterized by large pumice and black slag of the diameter of 20 - 30 cm. At the base incorporate numerous fragments of lava and limestone marl.
- **Tuffs yellow pumice clear:** The pumice tuff yellow light are very compact and generally fill paleo valleys excavated sediments Pliocene tuffs or yellow thick gray pumice.

With this acronym indicate levels of tuffs similar to the previous, but with included leucitic, to places with power exceeding 20 m. Surely the various lithological types from different eruptive vents. Typically these horizons are separated by paleo soils that sometimes show phenomena of cooking, which suggests that these levels may have set with phenomena similar to those of ignimbrites.

- **Tuffs:** This term indicates an alternation of layers of lapilli, pumice and cineritic separated by paleo soils blacks, each having a thickness varying from a minimum of 10 cm to a maximum of about 2 m. This alternating pyroclastic generally closes a series effusive local.

4.2 Spatial structural sub-surface

Based on the gravity data available in the literature it was outlined geometry of the structures that controlled sedimentation and volcanic activity during the Pliocene-Pleistocene area.

During the course of the Pleistocene the tectonic structures began to rally the regime undergoing subsidence of the central part. The continental basins linked to the sectors affected by such tectonic regime have developed just before and in conjunction Quaternary volcanic. Finally they set important features cross with anti-Apennines trend, visible in next figure.

The complex of Vulsini and Sabatini have developed coinciding with the two largest discontinuities observed that interrupt the Pliocene extensional structures and the dorsal midline

The transverse structure northernmost coincides with the alignment seismic identified to the north and east of Bolsena Lake [10]. During the course of volcanic Pleistocene, therefore, transverse structures trending anti-Apennines were activated, with a major role in guiding the ascent of

magmas potash. Next figure shows were major geological structures of the area in question (sedimentary basins, volcanic-tectonic structures and structural highs supposed/known) on the basis of the information gravimetric; many of these show evidence on the surface or seismic.





Diagram of the main sedimentary basins of Pliocene and Pleistocene, from literature data and the interpretation of the gravity data. 1) Complex volcanic acids Pliocene and Pleistocene. 2) Facilities bordiere major sedimentary basins of Pliocene and Pleistocene, deduced by gravimetric discontinuity. 3) Main facilities and caldera crater. 4) Main lava domes.

4.3 Earthquake

This study evidence an increase of Earthquake activity for 2006 to 2014. In next figure we can see the number of Earthquake in Acquapendente, and table evidence the number of Earthquake event by year.




Number of Earthquake for 1990 to 2014

Year	N. of Events	Year	N. of Events
2014	29	2013	14
2012	50	2011	65
2010	84	2009	43
2008	75	2007	18
2006	32	2005	3
2004	4	2003	0
2002	0	2001	4
2000	7	1999	5
1998	0	1997	1
1996	3	1995	0
1994	9	1993	8
1992	39	1991	2
1990	4		

Number of Earthquake in Acquapendente by Year

The study evidence that the major number of Earthquake are production on deep from 5 to 10 Km. Next graphics evidence this.



Magnitudo/Deep for 1990 to 2014


4.4 Wind speed

In this section we describe the wind speed. In the above image you can see the map of wind speed insistent on the territory of Acquapendente, as seen from the map generated from Atlas wind [6]. The study of daily wind speed has allowed to estimate on a monthly basis throughout the period included in this study:

- a decrease in the average for the period 1990-2014 in the speed of the winds, with values ranging from 9 Km/h (in September), and 21 Km/h in the month of July;
- an increase with higher gusts, in high winds, especially in the period between the autumn and winter, with values between 25 km/h (in August) and 36 Km/h (November).

The study also shows that the month of September is the one with the trend to-

wards greater variation in the wind speed, as well it appears from the comparison chart, maximum and minimum wind speed for this study shown in Figure 12.



Wind Speed Atlant of Acquapendente [6]

5 Conclusion

We presented for the first time an analysis of geological and longterm temperature data directly obtained from Acquapendente local meteorological site, inside urban concentration and well above the inversion layer.

The data analysis above all those related to annual rainfall, to their regimen and the change of these indices over time indicates that the area is recorded increase the contribution of rainfall and its concentration in the most rainy periods.

The extremes of this trend over time if confirmed, would lead to a radical change in climate of the area that would no longer be characterized by the absence of the dry season, but rather by a rainy season with

heavy and often heavy rainfall and a dry season with characteristics similar to those of the dry seasons of the Mediterranean climate.

The inversion of Acquapendente arouses contrasting effects among citizens, as it is a phenomenon particularly popular in the middle of a hot summer when at least in the early morning you are able to get much lower temperatures and refreshing, while in winter it is seen as a phenomenon inconvenient dates the low temperatures and possible fog and frost that usually occur in parallel.

Of course it is undoubtedly a characteristic phenomenon, which has advantages such as the cool summer morning or the ability to create a favorable condition to the snow fall.

Given the geology and structure of rocks and outcrops in the area, plus the non-poor seismic activity in the area, went increasing from 2006 until the preparation of this study, as shown in the table in section 4.2 of this paper, it is recommended further deepen the geological knowledge of the area in order to prepare the necessary protections for the populations and infrastructure present.

6 Acknowledgments

We would like to thank Silvia Gargano for her helpful suggestions and support to made the paper more complete. The constructive comments are highly appreciated.




Figure 1: Identification of Acquapendente in the Viterbo's province

Table 1: Comparison of temperature on decadal scale

Month	Year	Temperatures Min - Max	Month	Year	Temperatures Min - Max	Month	Year	Temperatures Min - Max
January	2014	5 - 11	February	2014	6 - 13	March	2014	5 - 16
	2013	3 - 10		2013	1 - 8		2013	5 - 12
	2012	2 - 11		2012	0 - 4		2012	6 - 19
	2011	3 - 10		2011	3 - 12		2011	5 - 13
	2010	2 - 8		2010	4 - 10		2010	4 - 13
	2009	4 - 9		2009	3 - 10		2009	6 - 14
	2008	4 - 12		2008	3 - 12		2008	5 - 14
	2007	4 - 13		2007	5 - 14		2007	6 - 16
	2006	0 - 8		2006	2 - 11		2006	4 - 13
	2005	1 - 9		2005	0 - 8		2005	3 - 14
	2004	1 - 9		2004	3 - 12		2004	4 - 13
	2003	3 - 10		2003	-1 - 9		2003	3 - 16
	2002	0 - 10		2002	5 - 14		2002	6 - 18
	2001	6 - 11		2001	3 - 11		2001	8 - 16
	2000	1 - 9		2000	4 - 13		2000	6 - 14
	1999	1 - 10		1999	1 - 10		1999	5 - 14
	1998	4 - 11		1998	4 - 14		1998	4 - 13
	1997	4 - 11		1997	3 - 12		1997	4 - 15
	1996	3 - 10		1996	0 - 8		1996	2 - 10
	1995	1 - 9		1995	4 - 12		1995	3 - 12
April	1994	4 - 10	May	1994	3 - 11	June	1994	4 - 16
	1993	2 - 10		1993	-1 - 11		1993	2 - 12
	1992	2 - 10		1992	2 - 12		1992	4 - 14
	1991	1 - 10		1991	1 - 9		1991	6 - 16
	1990	1 - 10		1990	3 - 13		1990	4 - 14
	2014	9 - 18		2014	11 - 21		2014	16 - 27
	2013	9 - 18		2013	10 - 19		2013	15 - 26
	2012	8 - 17		2012	11 - 21		2012	17 - 30
	2011	9 - 19		2011	12 - 24		2011	16 - 27
	2010	7 - 18		2010	12 - 20		2010	15 - 25
	2009	10 - 18		2009	15 - 25		2009	18 - 26
	2008	7 - 17		2008	11 - 22		2008	18 - 26
	2007	9 - 22		2007	12 - 24		2007	16 - 24
	2006	8 - 19		2006	11 - 24		2006	13 - 27
	2005	6 - 17		2005	12 - 25		2005	15 - 28
	2004	7 - 17		2004	9 - 20		2004	14 - 26
	2003	6 - 18		2003	9 - 21		2003	18 - 33
	2002	7 - 20		2002	9 - 19		2002	13 - 25
	2001	5 - 16		2001	12 - 23		2001	14 - 27
	2000	10 - 17		2000	14 - 24		2000	14 - 29
July	1999	8 - 16	August	1999	14 - 23	September	1999	18 - 28
	1998	8 - 17		1998	13 - 22		1998	16 - 27
	1997	4 - 14		1997	11 - 23		1997	15 - 26
	1996	6 - 16		1996	10 - 20		1996	14 - 27
	1995	4 - 14		1995	8 - 20		1995	11 - 24
	1994	5 - 15		1994	10 - 21		1994	14 - 25
	1993	6 - 17		1993	10 - 23		1993	14 - 27
	1992	7 - 16		1992	11 - 23		1992	13 - 23
	1991	3 - 14		1991	6 - 16		1991	12 - 24
	1990	4 - 13		1990	10 - 21		1990	14 - 25
	2014	7 - 27		2014	17 - 28		2014	15 - 25
	2013	19 - 30		2013	19 - 31		2013	15 - 26
	2012	19 - 33		2012	20 - 34		2012	16 - 25
	2011	17 - 27		2011	18 - 31		2011	16 - 28
	2010	19 - 31		2010	17 - 29		2010	14 - 24
	2009	20 - 30		2009	20 - 32		2009	16 - 26
	2008	19 - 30		2008	19 - 32		2008	15 - 25
	2007	17 - 32		2007	17 - 30		2007	14 - 26
	2006	18 - 33		2006	17 - 28		2006	16 - 27
	2005	17 - 31		2005	16 - 28		2005	14 - 25
	2004	16 - 30		2004	17 - 30		2004	14 - 27
October	2003	18 - 34	November	2003	19 - 33	December	2003	14 - 27
	2002	16 - 29		2003	16 - 27		2003	13 - 23
	2001	17 - 31		2001	18 - 32		2001	12 - 23
	2000	16 - 30		2000	17 - 33		2000	14 - 26
	1999	20 - 29		1999	20 - 31		1999	16 - 26
	1998	19 - 31		1998	19 - 31		1998	14 - 24
	1997	15 - 29		1997	17 - 29		1997	15 - 27
	1996	16 - 29		1996	16 - 28		1996	11 - 21
	1995	17 - 30		1995	15 - 26		1995	11 - 21
	1994	18 - 31		1994	19 - 32		1994	15 - 25
	1993	15 - 28		1993	18 - 32		1993	14 - 24
	1992	16 - 29		1992	18 - 32		1992	16 - 26
	1991	16 - 30		1991	17 - 31		1991	15 - 25
	1990	15 - 29		1990	16 - 28		1990	13 - 24
	2014	13 - 22		2014	10 - 17		2014	6 - 12
	2013	13 - 22		2013	8 - 14		2013	4 - 11
	2012	12 - 22		2012	9 - 15		2012	2 - 9
	2011	11 - 21		2011	7 - 16		2011	5 - 13
	2010	10 - 19		2010	7 - 14		2010	6 - 4
	2009	10 - 19		2009	7 - 16		2009	5 - 10
	2008	12 - 22		2008	8 - 15		2008	4 - 9
	2007	11 - 21		2007	5 - 14		2007	2 - 10
	2006	12 - 23		2006	7 - 17		2006	5 - 13
	2005	10 - 20		2005	4 - 8		2005	2 - 9
	2004	13 - 22		2004	7 - 14		2004	5 - 12
	2003	10 - 19		2003	8 - 15		2003	3 - 10
	2002	10 - 20		2003	10 - 17		2003	6 - 11
	2001	13 - 24		2001	6 - 15		2001	1 - 8
	2000	12 - 20		2000	8 - 16		2000	5 - 13
	1999	13 - 21		1999	7 - 14		1999	5 - 10
	1998	11 - 19		1998	6 - 12		1998	2 - 9
	1997	10 - 19		1997	8 - 14		1997	5 - 11
	1996	9 - 18		1996	7 - 14		1996	3 - 8
	1995	9 - 20		1995	4 - 12		1995	5 - 10
	1994	10 - 19		1994	8 - 15		1994	4 - 11
	1993	11 - 18		1993	5 - 11		1993	3 - 11
	1992	12 - 19		1992	7 - 15		1992	3 - 10
	1991	9 - 17		1991	5 - 12		1991	-1 - 8
	1990	11 - 19		1990	5 - 13		1990	1 - 7

Table 2: Day Time and Night Time Variation

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2014	6	7	9	9	10	11	10	11	9	10	8	6
2013	7	7	7	9	9	11	12	12	11	9	6	8
2012	9	9	12	9	10	13	13	14	9	9	7	7
2011	7	9	9	10	12	11	11	13	12	10	10	7
2010	6	6	9	10	8	10	12	12	10	9	7	6
2009	6	8	9	8	10	8	10	11	10	9	8	5
2008	8	10	9	10	10	8	10	13	10	9	8	5
2007	9	10	10	13	10	12	15	13	12	10	9	8
2006	9	9	9	11	13	11	14	11	11	10	10	8
2005	8	9	11	11	13	11	15	12	11	9	10	8
2004	8	8	10	10	12	14	15	14	12	9	7	6
2003	7	10	13	13	14	11	16	15	12	9	7	7
2002	10	9	12	13	12	14	13	11	10	10	7	6
2001	6	9	8	11	11	13	14	11	11	11	9	8
2000	8	8	8	8	10	15	14	15	12	9	7	7
1999	8	8	9	9	9	10	10	12	9	8	7	6
1998	7	10	9	9	9	11	11	12	10	8	6	7
1997	7	9	11	10	12	11	13	12	12	9	6	6
1996	6	8	8	10	10	12	13	12	10	8	7	5
1995	8	8	10	9	11	13	13	11	10	11	8	6
1994	7	8	12	9	11	11	13	14	11	9	6	7
1993	6	5	7	11	16	21	21	24	19	14	7	7
1992	9	10	10	9	12	10	12	14	13	7	8	7
1991	9	8	9	10	10	12	14	14	11	9	8	9
1990	9	10	10	9	11	13	14	13	11	8	8	6
Average 1990-2014	7,47	8,48	9,50	9,96	11,05	12,19	13,10	12,97	11,11	9,30	7,72	6,60

Table 3: Rain/Year [mm]



Year		Jan	Feb	Mar	Apr	May	Jun	Jul	Ago	Sep	Oct	Nov	Dec	Tot	% Year
2014	82	95,00	83,00	95,00	95,00	60,00	85,00	90,00	55,00	85,00	80,00	180,00	235,00	1238,00	5,78%
2013	128	185,00	155,00	140,00	35,00	80,00	155,00	255,00	15,00	50,00	65,00	95,00	25,00	1255,00	5,86%
2012	58	80,43	75,00	55,00	120,00	80,00	15,00	5,00	35,00	125,00	115,00	95,00	145,00	945,43	4,42%
2011	86	40,00	35,00	125,00	40,00	144,00	140,00	150,00	15,00	35,00	45,00	35,00	55,00	859,00	4,01%
2010	49	48,00	67,00	35,00	67,00	96,00	44,00	10,00	25,00	5,00	47,00	107,00	53,00	604,00	2,82%
2009	96	44,30	95,10	105,40	76,50	60,60	154,50	92,30	140,00	264,00	190,00	146,00	45,00	1413,70	6,61%
2008	98	71,00	53,50	93,00	104,00	146,20	81,90	32,80	99,70	51,40	146,40	285,90	16,00	1181,80	5,52%
2007	56	47,40	74,50	120,90	50,10	47,20	60,00	57,60	58,50	48,50	68,50	24,70	18,90	676,80	3,16%
2006	82	232,20	124,80	94,00	47,80	19,90	10,90	110,10	123,20	98,10	43,20	77,20	3,00	985,30	4,60%
2005	76	44,40	38,20	51,70	91,70	93,40	54,80	24,60	130,70	202,60	101,30	59,30	17,20	909,90	4,25%
2004	84	41,70	123,50	97,00	210,40	50,60	1,50	23,10	35,00	63,00	101,60	212,00	53,60	1013,00	4,73%
2003	63	84,20	24,40	17,00	23,20	140,10	57,10	53,90	25,70	95,40	94,40	115,80	20,80	752,00	3,51%
2002	64	32,00	37,70	11,40	43,50	63,20	34,10	76,20	110,80	132,30	18,20	82,70	123,20	765,30	3,58%
2001	43	142,20	34,90	83,30	74,20	27,30	17,00	20,60	8,00	22,50	11,60	62,60	13,80	518,00	2,42 %
2000	57	15,00	7,00	38,00	30,00	98,00	37,60	59,10	57,20	26,00	125,40	108,80	84,60	686,70	3,21%
1999	58	20,00	15,00	70,00	60,00	55,00	85,00	85,00	35,00	100,00	45,00	75,00	50,00	695,00	3,25%
1998	47	30,00	20,00	20,00	100,00	35,00	60,00	5,00	60,00	55,00	90,00	45,00	45,00	565,00	2,64%
1997	41	50,00	30,00	25,00	65,00	10,00	50,00	20,00	45,00	10,00	90,00	55,00	45,00	495,00	2,31%
1996	88	65,00	115,00	30,00	115,00	145,00	80,00	30,00	65,00	147,00	58,00	132,00	70,00	1052,00	4,92%
1995	84	45,00	60,00	70,00	93,00	105,00	20,00	30,00	280,00	90,00	15,00	30,00	170,00	1008,00	4,71%
1994	50	37,00	30,00	5,00	163,00	20,00	85,00	45,00	10,00	62,00	50,00	40,00	50,00	597,00	2,79%
1993	50	10,00	5,00	25,00	75,00	160,00	15,00	-	20,00	100,00	70,00	80,00	45,00	605,00	2,83%
1992	78	50,00	15,00	85,00	100,00	65,00	115,00	30,00	30,00	65,00	242,00	50,00	85,00	932,00	4,35%
1991	69	10,00	55,00	45,00	110,00	110,00	30,00	60,00	20,00	110,00	140,00	128,00	5,00	823,00	3,85%
1990	69	15,00	16,00	25,00	140,00	90,00	10,00	45,00	85,00	35,00	175,00	115,00	75,00	826,00	3,86%
Average 1990-2014	70,25	61,39	55,58	62,47	85,21	80,06	59,94	56,41	63,35	83,11	89,10	97,48	61,96	21401,93	
														Year Average	856,08

Table 4: Wind Average[1990-2014]/Year on Km/h

Day	Jan	Feb	Mar	Apr	May	Jun	Jul	Ago	Sep	Oct	Nov	Dec	Average	Min	Max
1	29.09	26.35	24.45	26.38	20.10	23.64	21.59	23.43	26.91	18.43	23.17	27.09	24.22	18.43	29.09
2	23.18	24.57	27.23	24.33	21.33	21.24	23.14	23.33	23.96	18.83	24.70	22.61	23.20	18.83	27.23
3	26.32	22.13	26.50	22.54	25.86	22.45	22.61	23.61	22.91	20.00	22.57	24.82	23.53	20.00	26.50
4	26.95	24.74	27.64	26.79	25.29	21.77	23.04	23.91	24.46	22.39	24.17	22.18	24.45	21.77	27.64
5	24.09	25.26	29.10	27.04	19.23	21.82	22.86	25.09	23.87	22.74	22.91	20.82	23.73	19.23	29.10
6	21.77	26.74	24.90	22.83	23.95	22.57	22.52	23.71	22.25	20.96	21.77	24.45	23.20	20.96	26.74
7	19.27	24.00	27.41	24.63	23.24	21.95	24.65	22.26	24.00	24.29	26.65	28.09	24.20	19.27	28.09
8	19.86	26.43	29.38	22.00	20.50	22.14	25.26	23.86	22.82	22.21	23.57	26.18	23.69	19.86	29.38
9	18.55	25.61	25.71	25.22	20.00	22.81	22.83	22.29	26.13	19.88	21.22	28.71	23.25	18.55	28.71
10	18.59	24.91	26.14	26.30	19.65	22.95	23.52	24.55	22.83	20.71	23.65	26.68	23.37	18.59	26.68
11	21.13	25.86	26.76	27.09	18.43	23.00	24.83	24.52	22.95	19.67	25.38	20.86	23.37	18.43	27.09
12	23.26	24.57	22.00	26.70	21.60	21.24	23.52	25.75	28.62	21.17	22.48	21.50	23.53	21.17	28.62
13	21.68	23.35	23.64	24.22	21.10	21.52	22.43	22.29	24.50	21.91	23.41	23.24	22.77	21.10	24.50
14	22.30	21.39	20.41	23.91	22.50	21.76	22.82	24.04	26.50	22.39	25.50	26.10	23.30	20.41	26.50
15	25.09	27.59	21.55	24.04	22.71	23.67	25.27	22.88	23.26	23.22	26.27	24.86	24.20	21.55	27.59
16	23.17	28.77	24.68	25.48	21.70	22.71	24.77	22.24	23.48	22.43	31.77	23.77	24.58	21.70	31.77
17	23.04	25.96	23.82	23.70	23.05	22.64	24.39	22.54	21.00	22.74	26.27	23.91	23.59	21.00	26.27
18	21.52	25.82	24.14	24.61	26.25	21.52	23.22	20.71	21.17	20.83	23.77	22.00	22.96	20.71	26.25
19	22.82	25.87	25.86	23.52	26.05	22.00	23.21	22.25	20.32	22.39	23.64	21.77	23.31	20.32	26.05
20	21.86	23.61	22.18	21.30	22.95	23.38	24.30	22.17	22.54	22.52	25.36	24.27	23.04	21.30	25.36
21	24.36	22.39	23.77	24.95	21.65	21.23	25.30	23.50	24.17	26.96	23.91	23.35	23.80	21.23	26.96
22	25.10	24.36	24.64	21.00	21.15	26.36	23.30	23.33	20.52	21.70	22.00	23.83	23.02	20.52	26.36
23	25.22	26.27	26.78	21.95	23.76	22.48	23.48	23.09	23.04	24.09	21.59	27.39	24.10	21.59	27.39
24	28.74	24.05	25.43	21.17	22.80	21.48	27.30	21.26	23.71	22.57	21.95	28.45	24.08	21.17	28.74
25	29.48	23.00	27.00	22.61	22.62	22.48	23.17	22.48	23.00	20.65	20.14	27.32	23.66	20.14	29.48
26	26.70	23.27	26.48	22.96	21.60	22.17	22.74	21.52	25.57	22.29	21.68	28.65	23.80	21.52	28.65
27	23.43	23.45	28.65	21.78	22.00	22.70	23.09	22.54	23.04	20.26	23.24	24.52	23.23	20.26	28.65
28	21.22	27.59	26.13	18.41	21.45	23.65	23.52	24.86	21.91	21.78	22.77	26.35	23.30	18.41	27.59
29	26.48	20.20	23.78	20.86	24.14	23.04	23.61	24.22	22.22	22.55	21.73	25.00	23.15	20.20	26.48
30	26.04		25.08	20.95	21.62	21.96	23.35	24.77	21.95	22.74	22.82	23.36	23.15	20.95	26.04
31	25.52		28.33	18.50	21.24	20.00	25.00	25.83	12.00	23.77	27.25	26.64	23.10	12.00	28.33
Average	23.74	24.76	25.47	23.48	22.24	22.40	23.70	23.29	23.08	21.90	23.78	24.80	Average	12.00	31.77
Min	18.55	20.20	20.41	18.41	18.43	20.00	21.59	20.71	12.00	18.43	20.14	20.82	22.77		
Max	29.48	28.77	29.38	27.09	26.25	26.36	27.30	25.83	28.62	26.96	31.77	28.71	24.58		

Table 5: Trend of Temperature of Acquapendente in Jan-Feb-Mar and Jun-Jul-Aug

Year	Jan	Feb	Mar	Jun	Jul	Aug
2014	7.68	9.79	10.06	21.37	22.15	22.84
2013	6.06	4.66	8.58	20.78	24.50	25.19
2012	6.35	1.42	12.58	23.48	25.90	27.08
2011	6.53	7.23	8.54	21.93	22.11	23.69
2010	5.10	6.79	8.56	19.48	24.90	23.31
2009	6.48	6.21	10.08	21.62	25.03	25.95
2008	7.77	7.51	9.34	21.57	24.61	25.53
2007	8.71	9.59	10.71	22.09	24.74	24.15
2006	4.04	6.44	8.44	20.29	25.58	22.76
2005	4.76	3.99	8.76	21.44	24.26	21.89
2004	4.88	7.48	8.28	19.38	24.74	23.98
2003	6.42	3.94	9.47	25.33	26.27	28.27
2002	4.00	7.65	12.26	23.85	23.15	21.45
2001	8.47	7.13	11.74	20.43	23.96	25.06
2000	5.35	8.40	10.05	21.36	22.87	24.93
1999	5.73	5.51	9.37	23.22	24.48	25.52
1998	7.06	8.67	8.55	21.92	24.95	25.03
1997	7.43	7.57	9.31	20.18	22.13	23.13
1996	6.62	3.91	5.94	20.36	22.26	22.00
1995	4.79	8.13	7.45	17.10	23.82	20.84
1994	6.95	6.75	10.05	19.20	24.42	25.39
1993	5.88	5.41	7.14	20.53	21.60	24.65
1992	6.00	6.81	9.05	18.30	22.50	25.16
1991	5.62	5.13	10.96	18.03	23.13	23.71
1990	5.77	7.82	9.23	18.76	22.10	21.98
Average 1990-2014	6.18	6.56	9.38	20.88	23.85	24.14






Figure 4: Maximum Temperature (solid line) and Average Temperature Variations 1990-2014 (dotted line)




Figure 5: Minimum Temperature (solid line) and Average Temperature Variations 1990-2014 (dotted line)




Figure 6: Total Days of Rain by Year




Figure 7: Total rain [mm] by year




Figure 8: Total day of Fog by year




Figure 9: Humidity (solid line) and Average Humidity Variations 1990-2014 (dotted line)




Figure 10: Pressure (solid line) and Average Pressure Variations 1990-2014 (dotted line)




Figure 11: Dew Point (solid line) and Average Dew Point Variations 1990-2014 (dotted line)




Figure 12: Wind Speed (solid line) and Average Wind Speed Variations 1990-2014 (dotted line)




Figure 13: August Temperature (solid line) and Average August Temperature 1990-2014 (dotted line)




Figure 14: February Temperature (solid line) and Average February Temperature 1990-2014 (dotted line)




Figure 15: Summer Temperature (solid line) and Average Temperature Variations 1990-2014 (dotted line). Jun: blu, Jul: red Aug: green




Figure 16: Winner Temperature (solid line) and Average Temperature Variations 1990-2014 (dotted line) Jan: blu, Feb: red Mar: green

Figure 17: Total Rain (mm) by Month





Figure 18: Solar Irradiation Map






Figure 19: DNI Direct Normal Irradiance (Wh/m²/day)

References

- [1] **Tasselli, D. Ricci, S.**, “Progetto Stazioni Meteorologiche e Sismologiche”, TS Corporation Srl - Dipartimento Meteorologia e Climatologia, 2014.
- [2] **Ente per le Nuove Tecnologie, l'Energia e l'Ambiente**, “Tabella dei gradi/giorno dei Comuni italiani raggruppati per Regione e Provincia, Legge 26 agosto 1993, n. 412, allegato A”, 1 marzo 2011, p. 151.
- [3] **Prof. L. Carmignani, Prof. G. Cornamusini, Dott. I. Callegari**, “Mappa Geologica di Acquapendente (VT)”, Università di Siena, Dipartimento di Scienze della Terra - Centro di GeoTecnologie, Foglio 333.
- [4] **IlMeteo.it** “Archivio Dati Meteo Acquapendente (VT)”, IlMeteo.it.
- [5] **Eumetsat** “Archivio Dati Meteo Acquapendente (VT)”, Eumetsat.
- [6] **Atlantedelvento** “Archivio Dati Atlante Eolico”, <http://www.atlanteeolico.it>.
- [7] **ISPRA** “Mappa Idrogeologica di Acquapendente (VT)”, ISPRA.
- [8] **Della Vedova et al.** “Geothermal structure of Tyrrhenian Sea. Mar. Geol. 55, 271–289”, 1994
- [9] **Arisi Rota et Fichera** “Magnetic interpretation connected to Geomagnetic provinces: the Italian case history. 47th Meeting European Association of Exploration Geophysicists Proc.”, 1985
- [10] **Buonasorte et al.** “Tectonic structures and geometric setting of the Vulsini Volcanic Complex. Per. Mineral., 56: 123-136”, 1987a
- [11] **F. Barberi, Buonasorte G., R. Cioni, Fiordelisi A., Foresi L., S. Iaccarino, Laurenzi MA, Sbrana A., L. Vernia, Villa IM** “Plio-Pleistocene geological evolution of the geothermal area of Tuscany and Latium. Mem. Descr. Carta Geol. D'It. XLIX (1994), pp 77-134”
- [12] **Iaccarino et al.** “Osservazioni stratigrafiche sul bordo orientale del Bacino di Radicofani. Mem. Descr. Carta Geol. D'It. XLIX”, 1994
- [13] **European Commission** “Map of Solar Irradiation”, 2012